



Mitochondrial Research



Mito-TEMPO



Mdivi

References

1. Ghosh *et al.* (2010) Free Radic.Biol.Med. **49** 1674
2. McManus *et al.* (2011) J.Neurosci. **31** 15703
3. Wang *et al.* (2013) PLoS One **8** e75044
4. Liang *et al.* (2010) Free Radic.Biol.Med. **49** 1550
5. Dikalova *et al.* (2010) Circulation Res. **107** 106
6. Kline *et al.* (2008) Exp.Hematol.**36** 568
7. Balakrishnan *et al.* (2009) Blood **113** 149
8. Cassidy-Stone *et al.* (2008) Dev.Cell **14** 193
9. Lackner *et al.* (2010) Chem.Biol. **17** 578
10. Benz and McLaughlin (1983) Biophys.J. **41** 381
11. Fulda and Kroemer (2009) Drug Discovery Today **14** 885
12. Seaton *et al.* (1998) Brain Res. **809** 12
13. Merlini *et al.* (2008) PNAS **105** 5225
14. Chelli *et al.* (2004) Biochem.Pharmacol. **68** 125
15. Sutter *et al.* (2004) J.Hepatol. **40** 1212
16. Maaser *et al.* (2001) Br.J.Cancer **85** 1771
17. Nguyen *et al.* (2007) PNAS **104** 19512
18. Huang *et al.* (2009) Clin.Cancer Res. **15** 1126
19. Andreaux *et al.* (2013) Nature Rev.Drug Disc. **12** 465
20. Long *et al.* (2016) Cell **166** 424

Mitoquinone

A mitochondria-targeted antioxidant displaying cardio- and neuroprotective effects.^{1,2}
10-1363 5 mg / \$70.00, 25 mg / \$280.00

Mito-TEMPO

A mitochondria-targeted antioxidant targeting reactive oxygen species.^{3,4,5}
10-4100 5 mg / \$55.00, 25 mg / \$225.00

AT-101

A small molecule mimic of the BH3 domain of cellular Bcl-2 inhibitors which interferes with the function of Bcl-2-family anti-apoptotic proteins. Induces apoptosis in drug-resistant multiple myeloma cell lines. Apoptosis is activated by the mitochondrial pathway. Cell permeable.^{6,7}

10-1165 10 mg / \$116.00, 50 mg / \$481.00

Mdivi

Selective inhibitor of Dnm1 GTPase ($IC_{50}=1\text{-}10 \mu\text{M}$). Inhibits mitochondrial outer membrane permeabilization. Inhibits mitochondrial division and apoptosis. Cell permeable.^{8,9}
10-1179 5 mg / \$40.00, 25 mg / \$160.00

FCCP

FCCP is a potent and reversible inhibitor of mitochondrial oxidative phosphorylation. It depolarizes mitochondrial membrane potential and induces apoptosis¹⁰
10-1182 10 mg / \$75.00, 50 mg / \$300.00

Betulinic Acid

Induces mitochondrial permeability pore opening leading to apoptosis in a variety of cell lines.¹¹

10-2278 10 mg / \$50.00, 50 mg / \$193.00

Cyclosporine A

Inhibits mitochondrial permeability pore opening inhibiting apoptosis.^{12,13}
10-1119 100 mg / \$50.00, 500 mg / \$200.00

PK-11195

PK-11195 is a selective and potent peripheral benzodiazepine receptor, PBR (or TSPO, translocator protein) antagonist. PK11195 is also able to induce apoptosis in a variety of cell lines.^{14,15,16}

10-2221 5 mg / \$55.00, 25 mg / \$220.00

Obatoclax

Bcl-2 antagonist that overcomes Bcl-2, Bcl-xL, Bcl-w and Mcl-1-mediated resistance to BAX or BAK.^{17,18}

10-2660 5 mg / \$70.00, 25 mg / \$276.00

M-1

Enhances mitochondrial fusion without interfering with endoplasmic reticulum and lysosome morphology.¹⁹

10-1598 5 mg / \$48.00, 25 mg / \$116.00

N-Oleoylphenylalanine

N-Oleoylphenylalanine is an endogenous uncoupler of UCP1-independent respiration.²⁰
10-1602 5 mg / \$50.00, 25 mg / \$200.00