MYC

The *MYC* family of proto-oncogenes encodes for three MYC oncoproteins: c-MYC, N-MYC and L-MYC.^{1,2} c-MYC is ubiquitous in both fetal and adult proliferating cells, while N- and L-MYC are more limited. These proteins are transcription factors that activate a large number of genes – 15% of all genes are believed to be regulated by MYC proteins.³ Cellular processes that are regulated by MYC include cell cycle, apoptosis, cell growth, differentiation, and metabolism.^{4,5} Because of its importance in these areas, >50% of solid tumors and leukemia feature deregulated MYC ^{6,7} and is, thus, an area of intense therapeutic interest.⁸

BI 2536

BI 2536 is a dual Plk/BRD4 inhibitor that destabilizes MYC via inhibiting the deactivation of ubiquitin E3 ligase Fbw7 by Plk1.⁹

Product No: 10-4121 5 mg/ 25 mg/__

P22077

An inhibitor of USP7 (HAUSP) that destabilizes N-MYC protein levels in a panel of neuroblastoma cell lines. $^{10}\,$

Product No: 10-4125 10 mg/ 50 mg/

Rapamycin

Rapamycin limited expression of c-MYC in granulocytes by inhibiting c-MYC mRNA translation.¹¹

Product No: 10-1104

10 mg/ 50 mg/__

MK-2206

MK2206 is a potent and highly specific allosteric pan-Akt inhibitor. Akt has been shown to act synergistically with c-MYC to activate tRNA synthesis and ribosome biogenesis in lymphomas.^{12,13}

Product No: 10-4801 10058-F4

<u>5 mg/ 25 mg/</u>

5 mg/ 25 mg/

10058-F4 disables c-MYC by inhibiting formation of the c-MYC-MAX complex.14

Product No: 10-1466

JQ1

BET bromodomain inhibitor that selectively downregulates MYC and MYCdependent target genes.¹⁵

Product No: 10-1361

THZ1

5 mg/ 25 mg/___

THZ1 is a potent covalent inhibitor of CDK7 that decreases STAT3 chromatin binding and MYC expression.¹⁶

Product No: 10-4120 5 mg/ 25 mg/

Stauprimide

Stauprimide selectively suppresses MYC transcription in a variety of cancer cell *in vitro* and inhibits tumor growth in mouse models.¹⁷

Product No: 10-1189 1 mg/ 5 mg/

Alisertib

Alisertib is a potent and selective inhibitor of Aurora A kinase (AurA). It disrupts the AurA/N-MYC complex leading to destabilization of N-MYC.¹⁸

Product N0: 10-4124 5 mg/ 25 mg/

10074-G5

Inhibits c-Myc by disrupting c-Myc/Max heterodimer formation and inhibiting its transcriptional activity. $^{19}\,$

Product N0: 10-5627 5 mg/ 25 mg/ I-BET762

BET bromodomain inhibitor that downregulates Myc transcription.²⁰

Product N0: 10-4122

5 mg/ 25 mg/

F CUS BIOMOLECULES



Alisertib

REFERENCES

- 1. Adhikary and Eilers (2005), Nat.Rev.Mol.Cell Biol. 6 635
- 2. Dang (2012), Cell 149 22
- 3. Dang et al. (2006), Semin.Cancer Biol. 16 253
- 4. Bretones et al. (2015), Biochim.Biophys. Acta 1849 506
- 5. Miller et al. (2012), Clin.Cancer Res. 18 5546
- 6. Vita and Hendriksson (2006), Semin.Cancer Biol. 16 318
- 7. Delgado and Leon (2010), Genes Cancer 1 605
- 8. Chen et al. (2018), Signal Transduct. Target Ther. 3 5
- 9. Xiao et al. (2016). Mol.Cell 64 493
- 10. Tavana et al. (2016) Nat.Med. 22 1180
- 11. Wall et al. (2008). Blood 112 2305
- 12. Chan et al. (2011), Sci.Signal. 4 ra56
- 13. Devlin et al. (2013), FEBS J. 280 5307
- 14. Yin et al. (2003), Oncogene 22 6151
- 15. Delmore et al. (2011), Cell 146 904
- 16. Cayrol et al. (2017), Nat. Commun. 8 14290
- 17. Bouvard et al. (2017), PNAS 114 3497
- 18. Richards et al. (2016), PNAS 113 13726
- 19. Follis et al. (2008), Chem. Biol. 15 1149
- 20. Delmore et al. (2011), Cell 146 904

400 Davis Dr. Suite 600 Plymouth Meeting, PA 19462 610-994-1134 sales@focusbiomolecules.com focusbiomolecules.com