

## Catalog # 10-1107 ONO-RS-082

CAS# 99754-06-0
2-(4-Pentylcinnamoyl)amino-4-chlorobenzoic acid; PACA
Lot # FBA6278

$$\bigcap_{\mathsf{N}} \bigcap_{\mathsf{CO}_2\mathsf{H}}^{\mathsf{CI}}$$

Reversible inhibitor of  $Ca^{2+}$ -independent phospholipase  $A_2$  (PLA<sub>2</sub>),  $IC_{50}$ =7  $\mu$ M for guinea pig lung PLA2.<sup>1</sup> Inhibits epinephrine-induced thromboxane production in platelets.<sup>1</sup> Inhibition of PLA<sub>2</sub> by ONO-RS-032 disrupts endosome tubule formation and maintenance of the Golgi complex.<sup>2-4</sup> Prevents Xenopus oocyte maturation at stage V.<sup>5</sup>

- 1) Banga et al. (1986), Activation of phospholipases A and C in human platelets exposed to epinephrine: role of glycoproteins Ilb/Illa and dual role of epinephrine; Proc.Natl.Acad.Sci.USA 83 9197
- 2) De Figueiredo *et al.* (2001), *Inhibition of transferrin recycling and endosome tubulation by phospholipase A2 antagonists*; J.Biol.Chem. **276** 47361
- 3) Schmidt et al. (2010), A role for phospholipase A2 activity in membrane tubule formation and TGN trafficking; Traffic 11 1530
- Bechler and Brown (2014), Gβ1γ2 activates phospholipase A2-dependent Golgi membrane tubule formation;
   Front.Cell.Dev.Biol. 28 0004
- 5) Islam et al. (2005), The distinct stage-specific effects of 2-(p-amylcinnamoyl)amino-4-chlorobenzoic acid on the activation of MAP kinase and Cdc2 kinase in Xenopus oocyte maturation; Cell Signal. **17** 507

## **PHYSICAL DATA**

NMR: (Conforms)

Solubility: Soluble in DMSO (25 mg/ml); ethanol (10 mg/mL)

Physical Description: White to off-white solid

Storage and Stability: Store as supplied at -20°C for up to 2 years from the date of purchase. Store DMSO or ethanol

solutions at -20°C for up to 3 months.

Materials provided by Focus Biomolecules are for laboratory research use only and are not intended for human or veterinary applications.