

Catalog # 10-1211 ODQ

CAS# 41443-28-1 1H-[1,2,4]Oxadiazolo[4,3-a]quinoxalin-1-one Lot # X101781

A potent and selective inhibitor of soluble guanylyl cyclase (sGC), $IC_{50} = 20 \text{ nM}$).¹ ODQ acts via competition with NO for the heme site of sGC where it binds irreversibly.² ODQ does not inhibit NO-mediated macrophage toxicity, an activity that is unrelated to cGMP nor does it inhibit particulate GC.¹ ODQ is an extremely useful tool to explore the involvement of the NO-cGMP pathway in cellular signaling and physiologic processes.³⁻⁵

- 1) Garthwaite et al. (1995), Potent and Selective Inhibition of Nitric Oxide-sensitive Guanylyl Cyclase by 1H-[1,2,4]Oxadiazolo[4,3-a]quinoxalin-1-one; Mol. Pharmacol. **48** 184
- 2) Schrammel et al. (1996), Characterization of 1H-[1,2,4]oxadiazolo[4,3-a]quinoxalin-1-one as a hemesite inhibitor of nitric oxide-sensitive guanylyl cyclase; Mol. Pharmacol. **50** 1
- 3) Estevez et al. (1998), Nitric oxide-dependent production of cGMP supports the survival of rat embryonic motor neurons cultured with brain-derived neurotrophic factor; J. Neurosci. **18** 3708
- 4) Vandecasteele et al. (1998), Role of the NO-cGMP in the muscarinic regulation of the L-type Ca²⁺ current in human atrial myocytes; J. Physiol. **506** 653
- 5) Martins-Pinge et al. (1999), Nitric oxide-dependent guanylyl cyclase participates in the glutamatergic neurotransmission within the rostral ventrolateral medulla of awake rats; Hypertension **34** 748

PHYSICAL DATA

Molecular Weight: 187.15 Molecular Formula: $C_9H_5N_3O_2$

Purity: >98% by HPLC

NMR: (Conforms)

Solubility: DMSO (40 mg/ml) Physical Description: Pale-yellow solid

Storage and Stability: Store as supplied at -20°C for up to 2 years from the date of purchase.

Solutions in DMSO may be stored at -20°C for up to 3 months