


## Catalog # 10-1281 Ribavirin

CAS# 36791-04-5 1-(β-D-Ribofuranosyl)-1H-1,2,4-triazole-3-carboxamide Lot # FBS1030



Ribavirin is a clinically useful antiviral medication for Hepatitis C, viral hemorrhagic fevers and other RNA and DNA viruses. The exact mechanism of its antiviral activity is uncertain. Various proposals include inhibition of viral polymerase<sup>1</sup>, RNA mutagenesis<sup>2</sup>, and inosine monophosphate dehydrogenase (IMPDH) inhibition<sup>3</sup>. Sensitizes tumor cells to anticancer agents 5-fluorouracil<sup>5</sup> and doxorubicin<sup>6</sup> via inhibition of eIF4E. Displays efficacy against atypical teratoid/rhabdoid tumors<sup>7</sup> and inhibits glioma cell growth<sup>8</sup>.

- 1) Bougie and Bisaillon (2003), Initial binding of the broad spectrum antiviral nucleoside ribavirin to the hepatitis C virus RNA polymerase; J. Biol. Chem. **278** 52471
- 2) Crotty et al. (2000), The broad-spectrum antiviral ribonucleoside ribavirin is an RNA virus mutagen; Nat.Med. 6 1375
- 3) Zhou et al. (2003), The effect of ribavirin and IMPDH inhibitos on hepatitis C virus subgenomic replicon RNA; Virology 310 333
- 4) Couee and Tipton (1990), Inhibition of ox brain glutamate by perphenazine; Biochem. Pharmacol. 39 1167
- 5) Hu et al. (2019), Ribavirin sensitizes nasopharyngeal carcinoma to 5-fluorouracil through suppressing 5-fluorouracil-induced ERK-dependentelF4E activation; Biochem. Biophys. Res. Commun. **513** 862
- 6) Tan et al. (2018), Ribavirin augments doxorubicin's efficicacy in human hepatocellular carcinoma through inhibiting doxorubicin-induced elF4E activation; J. Biochem. Mol. Toxicol. **32(1)** doi: 10.1002
- 7) Casaos et al. (2018), Ribavirin as a potential therapeutic for atypical teratoid/rhabdoid tumors; Oncotarget; 9 8054
- 8) Volpin et al. (2017) Use of an anti-viral drug, Ribavirin, as an anti-glioblastoma therapeutic; Oncogene, 36 3037

## PHYSICAL DATA

Molecular Weight:	244.21
Molecular Formula:	C <sub>8</sub> H <sub>12</sub> N <sub>4</sub> O <sub>5</sub>
Purity:	>98% by TLC
	NMR: (Conforms)
Solubility:	DMSO (25 mg/mL) and water (25mg/mL)
Physical Description:	White solid
Storage and Stability:	Store as supplied at -20°C for up to 1 year from the date of purchase. Solutions in
	DMSO or distilled water may be stored at -20°C for up to 3 months.

Materials provided by Focus Biomolecules are for laboratory research use only and are not intended for human or veterinary applications.

Focus Biomolecules LLC 400 Davis Drive, Suite 600 Plymouth Meeting PA 19462 www.focusbiomolecules.com