


Catalog # 10-1514

Salermide

CAS# 1105698-15-4

N-[3-[[[(2-Hydroxy-1-naphthalenyl)methylene]amino]phenyl]-α-methylbenzeneacetamide

Lot # S103093



SIRT1 and SIRT2 inhibitor. Exhibits a stronger inhibitory effect on SIRT2 than on SIRT1 *in vitro*. Induces the reactivation of proapoptotic genes repressed by SIRT1 and causes massive apoptosis in cancer cells within 24 hours.¹ Displays protective effect in nematodes expressing muscular dystrophy protein.² Promotes neuronal apoptosis confirming the protective effect of SIRT1.³

- 1) Lara *et al.* (2009), *Salermide, a Sirtuin inhibitor with a strong cancer-specific proapoptotic effect*; *Oncogene*, **28** 781
- 2) Pasco *et al.* (2010), *Characterization of sirtuin inhibitors in nematodes expressing a muscular dystrophy protein reveals muscle cell and behavioral protection by specific sirtinol analogues*; *J. Med. Chem.*, **53** 1407
- 3) Zhao *et al.* (2012), *Interactions between SIRT1 and MAPK/ERK regulate neuronal apoptosis induced by traumatic brain injury in vitro and in vivo*; *Exp. Neurol.*, **237** 489

PHYSICAL DATA

Molecular Weight:	394.47
Molecular Formula:	C ₂₆ H ₂₂ N ₂ O ₂
Purity:	98% by TLC
	NMR: (Conforms)
Solubility:	DMSO (up to 40 mg/ml), DMF (up to 30 mg/ml), or Ethanol (up to 10 mg/ml with warming)
Physical Description:	Yellow solid
Storage and Stability:	Store as supplied, desiccated at -20°C for up to 1 year from the date of purchase. Solutions in DMSO, DMF or ethanol may be stored at -20°C for up to 1 month.

Materials provided by Focus Biomolecules are for laboratory research use only and are not intended for human or veterinary applications.