


Catalog # 10-1583

Endosidin-2

CAS# 1839524-44-5

(E)-3-Fluoro-N'-(4-hydroxy-3-iodo-5-methoxybenzylidene)benzohydrazide
ES2

Lot # S103007



Endosidin-2 binds to the EXO70 subunit of the exocyst complex which results in inhibition of exocytosis and endosomal recycling in both plant and human cells and enhancement of plant vacuolar trafficking. EXO70 is a component of the octameric exocyst complex which tethers post-Golgi vesicles to the plasma membrane before SNARE-mediated membrane fusion². Endosidin-2 was previously identified as a plant endomembrane trafficking disruptor³. Inhibits intracellular glucose transporter, GLUT4 trafficking to the membrane and impairs glucose uptake in response to insulin⁴. It is a valuable new tool for the study of exocytosis.

- 1) Zhang *et al.* (2016), *Endosidin2 targets conserved exocyst complex subunit EXO70 to inhibit exocytosis*; Proc. Natl. Acad. Sci. USA, **113** E41
- 2) Novick *et al.* (1980), *Identification of 23 complementation groups required for post-translational events in the yeast secretory pathway*; Cell, **21** 205
- 3) Drakakaki *et al.* (2011), *Clusters of bioactive compounds target dynamic endomembrane networks in vivo*; Proc. Natl. Acad. Sci. USA, **108** 17850
- 4) Fujimoto *et al.* (2019), *The Exocyst Complex Regulates Insulin-Stimulated Glucose Uptake of Skeletal Muscle Cells*; Am. J. Physiol. Endocrinol. Metab. **317(6)** E957

PHYSICAL DATA

Molecular Weight:	414.17
Molecular Formula:	C ₁₅ H ₁₂ FIN ₂ O ₃
Purity:	98% by TLC
	NMR: (Conforms)
Solubility:	DMSO (at least 40 mg/ml), DMF (up to 35 mg/ml) or Ethanol (up to 10 mg/ml)
Physical Description:	Off-white solid
Storage and Stability:	Store as supplied at -20°C for up to 2 years from the date of purchase. Solutions in DMSO, DMF, or ethanol may be stored at -20°C for up to 1 month.

Materials provided by Focus Biomolecules are for laboratory research use only and are not intended for human or veterinary applications.