

Catalog # 10-1601 17-PAAG

17-(Propargylamino)-17-desmethoxygeldanamycin Lot # S103106

Semi-synthetic analog of geldanamycin which possesses an acetylenic side chain for coupling to other probes, fluorophores and bioactive molecules such as steroids via click chemistry. Chemical modifications at C17 are well known not to interfere with binding of geldanamycin to HSP90 client proteins. ²⁻⁴

- 1) Hendricks et al. (2013), Synthesis and preliminary evaluation steroidal anti-estrogen-geldanamycin conjugates; Bioorg. Med. Chem. Lett., **23** 3635
- 2) Llauger-Bufi et al. (2003), Synthesis of novel fluorescent probes for the molecular chaperone Hsp90; Bioorg. Med. Chem. Lett., **13** 3975
- 3) Kim et al. (2004), Development of a fluorescence polarization assay for the molecular chaperone Hsp90; J. Biomol. Screen., **9** 375
- 4) Austin et al. (2012), Fragment screening using capillary electrophoresis (CEfrag) for hit identification of heat shock protein 90 ATPase inhibitors; J. Biomol. Screen, **17** 868

PHYSICAL DATA

Molecular Weight: 583.67

 $\begin{array}{ll} \text{Molecular Formula:} & C_{31}H_{41}N_3O_8 \\ \text{Purity:} & 98\% \text{ by TLC} \end{array}$

NMR: (Conforms)

Solubility: Soluble in DMSO (up to at least 50 mg/ml) or in Ethanol (up to 5 mg/ml)

Physical Description: Violet solid

Storage and Stability: Store as supplied desiccated at -20°C for up to 1 year from the date of purchase.

Solutions in DMSO or ethanol may be stored at -20°C for up to 3 months.

Materials provided by Focus Biomolecules are for laboratory research use only and are not intended for human or veterinary applications.