

Catalog # 10-1608 Seriniquinone

22200-69-7 Dinaphthol[2,3-b:2',3'-d]thiophene-5,7,12,13-tetraone Lot # S103136

A new natural product isolated from a marine bacterium of the genus *Serinicoccus*. Displays selective toxicity against a distinct set of cell lines (within the NCI 60 panel) predominantly melanoma. Upon cellular treatment, seriniquinone was found to localize in the endoplasmic reticulum. Within 3 hours cells underwent autophagy followed by death via a caspase-9 apoptotic pathway. The target of seriniquinone was identified as the small protein, dermcidin as determined using an immunoaffinity approach.¹

 Trzoss et al. (2014), Seriniquinone, a selective anticancer agent, indices cell death by autophagocytosis, targeting the cancer-protective protein dermcidin; Proc. Natl. Acad. Sci. USA, 111 14687

PHYSICAL DATA

 $\begin{array}{lll} \mbox{Molecular Weight:} & 344.34 \\ \mbox{Molecular Formula:} & C_{20}\mbox{H}_8\mbox{O}_4\mbox{S} \\ \mbox{Purity:} & 98\% \ \mbox{by TLC} \\ \mbox{NMR: (Conforms)} \end{array}$

Solubility: Soluble in DMSO (up to 15 mg/ml with warming) or in DMF (up to 15 mg/ml)

Physical Description: Orange solid

Storage and Stability: Store as supplied desiccated at -20°C for up to 1 year from the date of purchase.

Solutions in DMSO or DMF may be stored at -20°C for up to 3 months.

Materials provided by Focus Biomolecules are for laboratory research use only and are not intended for human or veterinary applications.

Focus Biomolecules LLC 400 Davis Drive, Suite 600 Plymouth Meeting PA 19462 www.focusbiomolecules.com