

Catalog # 10-2449 Cyclothiazide

CAS# 2259-96-3

6-Chloro-3,4-dihydro-3-(5-norbornen-2-yl)-2*H*-1,2,4-benzothiadiazine-7-sulfonamide-1,1-dioxide LY-35483 Lot # FBS2022

Positive allosteric modulator of AMPA receptors acting at a site distinct from that of 2,3-benzodiazepines.^{1,2} Clinically useful diuretic and antihypertensive agent.³ Induces robust epileptiform activity, inducing seizures but without neuronal death.^{4,5} Can be used to produce a new animal model for epilepsy.⁵

- 1) Donevan and Rogawski (1998), Allosteric regulation of alpha-amino-3-hydroxy-5-methyl-4-isoxazole-propionate receptors by thiocyanate and cyclothiazide at a common modulatory site distinct from that of 2,3-benzodiazepines; Neuroscience **87** 615
- Desai et al. (1995), Cyclothiazide acts at a site on the alpha-amino-3-hydroxy-5-methyl-4-isoxazole propionic acid receptor complex that does not recognize competitive or noncompetitive AMPA receptor antagonists;
 J. Pharmacol. Exp. Ther. 272 38
- 3) Jeunemaitre *et al.* (1988), Long-term metabolic effects of spironolactone and thiazides combined with potassium-sparing agents for treatment of essential hypertension; Am. J. Cardiol. **62** 1072
- *4)* Qi *et al.* (2006), Cyclothiazide induces robust epileptiform activity in rat hippocampal neurons both in vitro and in vivo; J. Physiol. **571** 605
- 5) Kong et al. (2010), Cyclothiazide induces seizure behavior in freely moving rats; Brain Res. 1355 207

PHYSICAL DATA

Molecular Weight: 389.88

Molecular Formula: $C_{14}H_{16}CIN_3O_4S_2$ Purity: >98% by TLC NMR: (Conforms)

Solubility: DMSO (up to 35 mg/ml), or Ethanol (up to 9 mg/ml)

Physical Description: Off-white solid

Storage and Stability: Store as supplied at room temperature for up to 1 years from the date of purchase. Solutions in

DMSO or ethanol may be stored at -20°C for up to 3 months.

Materials provided by Focus Biomolecules are for laboratory research use only and are not intended for human or veterinary applications.