




Catalog # 10-2629

Monastrol

CAS# 254753-54-3

1,2,3,4-tetrahydro-4-(3-hydroxyphenyl)-6-methyl-2-thioxo-5-pyrimidinecarboxylic acid, ethyl ester

Lot # X101126



Potent, cell-permeable, small molecule inhibitor of mitosis. Monastrol-arrested cells are characterized by monopolar spindles.¹ Does not affect other motor proteins or tubulin. Specifically inhibits the mitotic molecular motor kinesin Eg5, a motor protein required for spindle formation ($IC_{50} = 14 \mu M$).² Mimics the effect of amyloid β on long term potentiation in a cellular model of learning and memory.³ Enhances regeneration of adult axons.⁴

- 1) Mayer *et al.* (1999), *Small molecule inhibitor of mitotic spindle bipolarity identified in a phenotype-based screen*; Science, **286** 971
- 2) Kapoor *et al.* (2000), *Probing spindle assembly mechanisms with monastrol, a small molecule inhibitor of mitotic kinesin, Eg5*; J. Cell Biol., **150** 975
- 3) Ari *et al.* (2014), *Alzheimer amyloid beta inhibition of Eg5/kinesin 5 reduces neurotrophin and/or reansmitter receptor function*; Neurobiol. Aging, **35** 1839
- 4) Lin *et al.* (2011), *Inhibition of Kinesin-5, a microtubule-based motor protein, as a strategy for enhancing regeneration of adult axons*; Traffic, **12** 269

PHYSICAL DATA

Molecular Weight:	292.35
Molecular Formula:	C ₁₄ H ₁₆ N ₂ O ₃ S
Purity:	98% by TLC
	NMR: (Conforms)
Solubility:	DMSO (up to 25 mg/ml) or Ethanol (up to 25 mg/ml)
Physical Description:	Beige solid
Storage and Stability:	Store as supplied, desiccated at -20°C for up to 1 year from the date of purchase. Solutions in DMSO or ethanol may be stored at -20°C for up to 1 month.

Materials provided by Focus Biomolecules are for laboratory research use only and are not intended for human or veterinary applications.

Focus Biomolecules LLC 400 Davis Drive, Suite 600 Plymouth Meeting PA 19462

www.focusbiomolecules.com