


Catalog # 10-2668 Diphenyleneiodonium CI

4673-26-1 DPI Lot # X101471



An irreversible inhibitor of iNOS and eNOS (IC₅₀s=50 and 100 nM respectively.¹ Binds to and inhibits the flavoprotein of neutrophil NADPH oxidase.² Novel agonist for GPR3.³ Inhibits NFκB activation induced by IL-1β.⁴ Prevents early alcohol-induced liver injury in rodent models.⁵

- 1) Stuehr et al. (1991), Inhibition of macrophage and endothelial cell nitric oxide synthase by diphenyleneiodonium and its analogs; FASEB J., 5 98
- Yea et al. (1990), Purification and some properties of the 45 kDa diphenylene iodonium-binding flavoprotein of neutrophil NADPH oxidase;
 Biochem. J., 265 95
- 3) Ye et al. (2014), Identification of a novel small-molecule agonist for human G protein-coupled receptor 3; J. Pharmacol. Exp. Ther., 349 437
- 4) Mendes et al. (2001), Diphenyleneiodonium inhibits NF-kappaB activation and iNOS expression induced by IL-1beta: involvement of reactive oxygen species; Mediators Inflamm., 10 209
- 5) Kono et al. (2001), Diphenyleneiodonium sulfate, an NADPH oxidase inhibitor, prevents early alcohol-induced liver injury in the rat, Am. J. Physiol. Gastrointest. Liver Physiol., **280** G1005

PHYSICAL DATA

Molecular Weight: 314.55

Molecular Formula: C₁₂H₈I ⋅ Cl

Purity: 98% by TLC

NMR: (Conforms)

Solubility: DMSO (up to 5 mg/ml with warming)

Physical Description: White solid

Storage and Stability: Store as supplied desiccated at -20°C for up to 1 year from the date of purchase. Solutions in

DMSO may be stored at -20°C for up to 2 months.

Materials provided by Focus Biomolecules are for laboratory research use only and are not intended for human or veterinary applications.