

Catalog # 10-2782 TUDCA

CAS# 14605-22-2 3α,7β-Dihydroxy-5β-cholan-24-oic acid N-(2-sulfoethyl)amide Tauroursodeoxycholic acid, sodium salt Lot # X106723



A classic inhibitor of endoplasmic reticulum (ER) stress.¹ Reduces ER stress and adipose tissue inflammation in a mouse model of high fat diet-induced obesity.² Blocks ER stress-induced NLRP3 inflammasome activation and hepatocyte death.³ Attenuates amyloid precursor protein processing and amyloid-β deposition in APP/PS1 mouse model.⁴

- 1) Xie et al. (2002), *Effects of tauroursodeoxycholic acid on endoplasmic reticulum stress-induced caspase-12 activation*; Hepatology, **36** 592
- 2) Chen et al. (2016), Chemical chaperones reduce ER stress and adipose tissue inflammation in high fat dietinduced mouse model of obesity; Sci.Rep., **6** 27486
- 3) Lebeaupin *et al.* (2015), *ER stress induces NLRP3 inflammasome activation and hepatocyte death*; Cell Death Dis., **6** e1879
- 4) Nunes et al. (2012), TUDCA, a bile acid, attenuates amyloid precursor protein processing and amyloid-β deposition in APP/PS1 mice; Mol.Neurobiol. 45 440

PHYSICAL DATA

Molecular Weight:	521.69
Molecular Formula:	$C_{26}H_{44}NO_6SNa$
Purity:	95% by TLC
	NMR: (Conforms)
Solubility:	Soluble in DMSO (up to 30 mg/ml) or in Ethanol (up to 20 mg/ml with warming)
Physical Description:	White solid
Storage and Stability:	Store as supplied desiccated at -20°C for up to 1 year from the date of purchase.
	Solutions in DMSO or ethanol may be stored at -20°C for up to 1 month.

Materials provided by Focus Biomolecules are for laboratory research use only and are not intended for human or veterinary applications.