

Catalog # 10-4102 Preladenant

CAS# 377727-87-2

4-(Furan-2-yl)-10-[2-[4-[4-(2-methoxyethoxy)phenyl]piperazin-1-yl]ethyl]-3,5,6,8,10,11-hexazatricyclo[7.3.0.0^{2,6}]dodeca-1(9),2,4,7,11-pentaen-7-amine; SCH 420814

Lot # FBS2013

Preladenant is a potent (Ki = 1.1nM) and selective (>1000-fold over other adenosine receptors) adenosine A_{2A} antagonist.¹ It has shown efficacy in rodent and primate models of Parkinsons disease without inducing dyskinesias and displays antidepressant effects.¹⁻³ It has been shown that activation of the adenosine A_{2A} receptor blocks the activation of immune cells and increases the number of regulatory T-cells.⁴⁻⁶ Currently in clinical trials as combination therapy with pembrolizumab.⁷

- 1) Hodgson et al. (2009), Characterization of the potent and selective A2A receptor antagonists preladenant and SCH 412348 [7-[2-[4-2,4-difluorophenyl]-1-piperazinyl]ethyl]-2-(2-furanyl)-7H-pyrazolo[4,3-e][1,2,4]triazolo[1,5-c]pyrimodin-5-amine] in rat models of movement disorders and depression; J.Pharmacol.Exp.Ther. **330** 294
- 2) Hodgson et al. (2010), Preladenant, a selective A(2A) receptor antagonist, is active in primate models of movement disorders; Exp.Neurol. **225** 384
- 3) Pinna et al. (2016), Antidyskinetic effect of A2A and 5HT1A/B receptor ligands in two animal models of Parkinson's disease; Mov.Disord. **31** 501
- 4) Beavis et al. (2013), Blockade of A2A receptors potently suppresses the metastasis of CD73+ tumors; Proc.Natl.Acad.Sci USA. **110** 14711
- 5) Hatfield and Sitkovsky (2016), A2A adenosine receptor antagonist to weaken the hypoxia-HIF-1a driven immunosuppression and improve immunotherapies of cancer; Curr.Opin.Pharmacol. **29** 90
- 6) Ohta et al. (2016), A metabolic immune checkpoint: adenosine in the tumor microenvironment; Front.Immunol. 7 1
- 7) NCT03099161

PHYSICAL DATA

Molecular Weight: 503.57 Molecular Formula: $C_{25}H_{29}N_9O_3$ Purity: >98% by HPLC NMR: (Conforms)

Solubility: DMSO (5 mg/ml with warming)

Physical Description: Beige solid

Storage and Stability: Store as supplied at -20°C for up to 1 year from the date of purchase. Solutions in

DMSO may be stored at -20°C for up to 1 month.

Materials provided by Focus Biomolecules are for laboratory research use only and are not intended for human or veterinary applications.