

Catalog # 10-4820 Idelalisib CAS# 870281-82-6

5-Fluoro-3-phenhyl-2-[(1S)-1-(7H-purin-6-ylamino)propyl]quinazolin-4-one; CAL-101 Lot # FBS1112

Idelalisib is a potent (IC $_{50}$ = 2.5nM) and selective (IC $_{50}$'s: PI3K $_{\alpha}$ = 820nM, PI3K $_{\beta}$ = 565nM, PI3K $_{\gamma}$ = 89nM) PI3K $_{\delta}$ inhibitor. Useful clinical agent for the treatment of various blood cancers. Idelalisib attenuates regulatory T cells (Treg) but not conventional T cells (Tconv) resulting in a significant increase in tumor-infiltrating antigen-specific CD8 T cells in a murine lung cancer model. Conversely, systemic PI3K $_{\delta}$ inactivation antagonized anti-CTLA-4 and anti-PD-L1 treatment. Others have found that Idelalisib minimally influenced rituximab- and obinutuzumab-mediated Ab-dependent cellular cytotoxicity in human lymphoma cells.

- 1) Herman et al. (2010) Phosphatidylinositol 2-kinase-δ inhibitor CAL-101 shows promising preclinical activity in chronic lymphocytic leukemia by antagonizing intrinsic and extrinsic cellular survival signals; Blood **116** 2078
- 2) Lannutti et al. (2011) CAL-101, a p110d selective phosphatidylinositol-3-kinase inhibitor for the treatment of B-cell malignancies, inhibits PI3K signaling and cellular viability; Blood **117** 591
- 3) Ahmad et al. (2017) Differential PI3Kd Signaling in CD4+ T-cell Subsets Enables Selective Targeting of T Regulatory Cells to Enhance Cancer Immunotherapy; Cancer Res. 77 1892
- 4) Lim et al. (2018) Phosphoinositide 3-kinase δ inhibition promotes antitumor responses but antagonizes checkpoint inhibitors; JCI Insight **3** e120626
- 5) Palazzo et al. (2018) The PI3Kd-Selective Inhibitor Idelalisib Minimally Interferes with Immune Effector Function Mediated by Rituximab or Obinutuzumab and Significantly Augment B Cell Depletion In Vivo; J.Immunol. **200** 2304

PHYSICAL DATA

Molecular Weight: 415.43

Molecular Formula: C₂₁H₁₈FN₇O

Purity: >98% by HPLC

NMR: (Conforms)

Solubility: DMSO (>25 mg/ml)

Physical Description: White solid

Storage and Stability: Store as supplied at -20°C for up to 1 year from the date of purchase. Solutions in

DMSO may be stored at -20°C for up to 1 month.

Materials provided by Focus Biomolecules are for laboratory research use only and are not intended for human or veterinary applications.