
Catalog #10-5244

BRM014

CAS 2270879-17-7

1-[3-(Difluoromethyl)isothiazol-5-yl]-3-[2-fluoro-5-(hydroxymethyl)pyridine-4-yl]urea; NVP-RXI570

Lot # E103775



SWI/SNF-related, matrix-associated, actin-dependent regulator of chromatin subfamily A member 2 (SMARCA2), also known as Brahma homologue (BRM), together with its close homologue Brahma-related gene 1 (BRG1), or SMARCA4, are ATPases of the large ATP-dependent SWI/SNF chromatin-remodeling complexes involved in transcriptional regulation of gene expression. BRM014 is an allosteric dual inhibitor of BRM and BRG1 (IC_{50}<5 nM for both).[1] It downregulated BRM-dependent gene expression and displayed antiproliferative activity in a BRG1-mutant-lung-tumor xenograft model after oral administration. Inhibition of BRG1/BRM ATPase by BRM014 inhibits chromatin openness.[2] For example, it reduced DNA accessibility to core regulatory factors (RUNX1, LM01 and MEIS1) at transcription factor PU.1 sites in AML cells.[3] P583 cells formed markedly large spheroids when treated with BRM014.[4] Cell permeable and active *in vivo*.

1) Papillon *et al.* (2018) *Discovery of orally active inhibitors of brahma homolog (BRM)/SMARCA2 ATPase activity for the treatment of brahma related gene 1 (BRG1)/SMARCA4-mutant cancers;* J. Med. Chem. **61** 10155
2) Hargreaves *et al.* (2021) *Chromatin openness requires continuous SW1/SNF activity;* Nat. Genet. **53** 263
3) Chambers *et al.* (2023) *SW1/SNF Blockade Disrupts PU.1-Directed Enhancer Programs in Normal Hematopoietic Cells and Acute Myeloid Leukemia;* Cancer Res. **83** 983
4) Krieg *et al.* (2023) *Chemical inhibition of SWI/SNF induces transformed growth in non-mutant cells, mimicking the effects of a SMARC4 mutation associated with small cell carcinoma of the ovary, hypercalcemic type;* Gynecol. Oncol. Rep. **48 Suppl 1** S32

PHYSICAL DATA

Molecular Weight:	318.27
Molecular Formula:	$C_{11}H_9F_3N_4O_2S$
Purity:	>98% HPLC
	NMR: (Conforms)
Solubility:	DMSO (70 mg/ml)
Physical Description:	White solid
Storage and Stability:	Store as supplied at -20°C for up to 2 years from the date of purchase. Solutions in DMSO may be stored at -20°C for up to 3 months.

Materials provided by Focus Biomolecules are for laboratory research use only and are not intended for human or veterinary applications.

Focus Biomolecules LLC 400 Davis Drive, Suite 600 Plymouth Meeting PA 19462

www.focusbiomolecules.com