

## Catalog #10-4650 P505-15

CAS# 1370261-97-4

2-[[(1R,2S)-2-Aminocyclohexyl]amino]-4-[3-(triazol-2-yl)anilino]pyrimidine-5-carboxamide hydrochloride; PRT062607; BIIB-057

## Lot # FBS4025

P505-15 is a highly selective and potent (IC50 = 1-2 nM) inhibitor of spleen tyrosine kinase (Syk). It potently inhibited B cell antigen receptor-mediated B cell signaling/activation and basophil degranulation. It displayed anti-inflammatory activity in rodent models of rheumatoid arthritis. P505-15 displayed efficacy in vitro and in vivo in non-Hodgkin lymphoma and chronic lymphocytic leukemia models. It protected against toxin-induced hepatic fibrosis, associated hepatocellular injury, and hepatocarcinogenesis. P505-15 was also found to inhibit PINK1 (IC50 = 0.5-3  $\mu$ M) in HeLa cells and dopaminergic neurons.

- 1) Coffey et al. (2012), Specific inhibition of spleen tyrosine kinase suppresses leukocyte immune function and inflammation in animal models of rheumatoid arthritis; J. Pharmacol. Exp. Ther. **340** 350
- 2) Hoellenriegel et al. (2012), Selective, novel spleen tyrosine kinase (Syk) inhibitors suppress chronic lymphocytic leukemia B-cell activation and migration; Leukemia 26 1576
- 3) Spurgeon et al. (2013), The selective SYK inhibitor P505-15 (PRT062607) inhibits B cell signaling and function in vitro and in vivo and augments the activity of fludarabine in chronic lymphocytic leukemia; J. Pharmacol. Exp. Ther. **344** 378
- 4) Torres-Hernandex et al. (2019), Targeting SYK signaling in myeloid cells protects against liver fibrosis and hepatocarcinogenesis; Oncogene **38** 4512
- 5) Rasool *et al.* (2024), *Identification and structural characterization of small molecule inhibitors of PINK1*; Sci. Rep. **14** 7739

## PHYSICAL DATA

Molecular Weight: 429.91

Molecular Formula:  $C_{19}H_{23}N_9O \cdot HCI$ Purity: >98% (HPLC)

NMR: (Conforms)

Solubility: DMSO (25 mg/mL); Water (>25 mg/mL)

Physical Description: White solid

Storage and Stability: Store as supplied at -20°C for up to 2 years from the date of purchase. Solutions in

DMSO or water may be stored at -20°C for up to 3 months.

Materials provided by Focus Biomolecules are for laboratory research use only and are not intended for human or veterinary applications.

Focus Biomolecules LLC 400 Davis Drive, Suite 600 Plymouth Meeting PA 19462